

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### GLUCOPHAGE 1000 mg film-coated tablets metformin hydrochloride

**Read all of this leaflet carefully before you start taking this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

#### **In this leaflet:**

1. What Glucophage is and what it is used for
2. Before you take Glucophage
3. How to take Glucophage
4. Possible side effects
5. How to store Glucophage
6. Further information

## **1. WHAT GLUCOPHAGE IS AND WHAT IT IS USED FOR**

### **What Glucophage is**

Glucophage contains metformin, a medicine to treat diabetes. It belongs to a group of medicines called biguanides.

Insulin is a hormone produced by the pancreas that makes your body take in glucose (sugar) from the blood. Your body uses glucose to produce energy or stores it for future use.

If you have diabetes, your pancreas does not make enough insulin or your body is not able to use properly the insulin it produces. This leads to a high level of glucose in your blood. Glucophage helps to lower your blood glucose to as normal a level as possible.

If you are an overweight adult, taking Glucophage over a long period of time also helps to lower the risk of complications associated with diabetes. Glucophage is associated with either a stable body weight or modest weight loss.

### **What Glucophage is used for**

Glucophage is used to treat patients with type 2 diabetes (also called 'non-insulin dependent diabetes') when diet and exercise alone have not been enough to control your blood glucose levels. It is used particularly in overweight patients.

Adults can take Glucophage on its own or together with other medicines to treat diabetes (medicines taken by mouth or insulin).

Children 10 years and over and adolescents can take Glucophage on its own or together with insulin.

## **2. BEFORE YOU TAKE GLUCOPHAGE**

### **Do not take Glucophage**

- if you are allergic (hypersensitive) to metformin or any of the other ingredients of this medicine (see 'What Glucophage contains' in section 6)
- if you have kidney or liver problems

- if you have uncontrolled diabetes, with e.g. severe hyperglycaemia (high blood glucose), nausea, vomiting, dehydration, rapid weight loss or ketoacidosis. Ketoacidosis is a condition in which substances called 'ketone bodies' accumulate in the blood and which can lead to diabetic pre-coma. Symptoms include stomach pain, fast and deep breathing, sleepiness or unusual fruity odour of the breath
- if you lost too much water from your body (dehydration), such as due to long-lasting or severe diarrhoea, or if you have vomited several times in a row. Dehydration may lead to kidney problems, which can put you at risk for lactic acidosis (see 'Take special care with Glucophage' below).
- if you have a severe infection, such as an infection affecting your lung or bronchial system or your kidney. Severe infections may lead to kidney problems, which can put you at risk for lactic acidosis (see 'Take special care with Glucophage' below).
- if you are treated for heart failure or have recently had a heart attack, have severe problems with your circulation (such as shock) or have breathing difficulties. This may lead to a lack in oxygen supply to tissue which can put you at risk for lactic acidosis (see 'Take special care with Glucophage' below)
- if you drink a lot of alcohol

If any of the above applies to you, talk to your doctor, before you start taking this medicine.

Make sure you ask your doctor for advice, if

- you need to have an examination such as X-ray or scan involving the injection of contrast medicines that contain iodine into your bloodstream
- you need to have major surgery

You must stop taking Glucophage for a certain period of time before and after the examination or the surgery. Your doctor will decide whether you need any other treatment for this time. It is important that you follow your doctor's instructions precisely.

### **Take special care with Glucophage**

#### **Please note the following particular risk of lactic acidosis.**

Glucophage may cause a very rare, but serious complication called lactic acidosis, particularly if your kidneys are not working properly. The risk of lactic acidosis is also increased with uncontrolled diabetes, prolonged fasting or alcohol intake. Symptoms of lactic acidosis are vomiting, bellyache (abdominal pain) with muscle cramps, a general feeling of not being well with severe tiredness, and difficulty in breathing. **If this happens to you, you may need immediate hospital treatment, as lactic acidosis may lead to coma. Stop taking Glucophage immediately and contact a doctor or the nearest hospital straight away.**

Glucophage on its own does not cause hypoglycaemia (a blood glucose level which is too low). However, if you take Glucophage together with other medicines to treat diabetes that can cause hypoglycaemia (such as sulphonylureas, insulin, meglitinides), there is a risk of hypoglycaemia. If you experience symptoms of hypoglycaemia such as weakness, dizziness, increased sweating, fast heart beating, visions disorders or difficulty in concentration, it usually helps to eat or drink something containing sugar.

### **Taking other medicines**

If you need to have an injection of contrast medicines that contain iodine into your bloodstream, for example for examinations such as X-ray or scan, **you must stop taking Glucophage for a certain period of time before and after the examination** (see 'Make sure you ask your doctor for advice' above).

Tell your doctor if you take any of the following medicines and Glucophage at the same time. You may need more frequent blood glucose tests or your doctor may adjust the dosage of Glucophage:

- diuretics (used to remove water from the body by making more urine).
- beta-2 agonists such as salbutamol or terbutaline (used to treat asthma)

- corticosteroids (used to treat a variety of conditions, such as severe inflammation of the skin or in asthma)
- other medicines used to treat diabetes

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

### **Taking Glucophage with food and drink**

Do not drink alcohol when you take this medicine. Alcohol may increase the risk of lactic acidosis especially if you have liver problems or if you are undernourished. This also applies to medicines that contain alcohol.

### **Pregnancy and breast-feeding**

During pregnancy, you need insulin to treat your diabetes. Tell your doctor if you are, you think you might be or are planning to become pregnant, so that he or she may change your treatment. This medicine is not recommended if you are breast-feeding or if you are planning to breast-feed your baby.

### **Driving and using machines**

Glucophage on its own does not cause hypoglycaemia (a blood glucose level which is too low). This means that it will not affect your ability to drive or use machines.

However, take special care if you take Glucophage together with other medicines to treat diabetes that can cause hypoglycaemia (such as sulphonylureas, insulin, meglitinides). Symptoms of hypoglycaemia include weakness, dizziness, increased sweating, fast heart beat, vision disorders or difficulty in concentration. Do not drive or use machines if you start to feel these symptoms.

## **3. HOW TO TAKE GLUCOPHAGE**

Always take Glucophage exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Glucophage cannot replace the benefits of a healthy lifestyle. Continue to follow any advice about diet that your doctor has given you and get some regular exercise.

### **Usual dose**

Children 10 years and over and adolescents usually start with 500 mg or 850 mg Glucophage once a day. The maximum daily dose is 2000 mg taken as 2 or 3 divided doses. Treatment of children between 10 and 12 years of age is only recommended on specific advice from your doctor, as experience in this age group is limited.

Adults usually start with 500 mg or 850 mg Glucophage two or three times a day. The maximum daily dose is 3000 mg taken as 3 divided doses.

If you take insulin too, your doctor will tell you how to start Glucophage.

### **Monitoring**

- Your doctor will perform regular blood glucose tests and will adapt your dose of Glucophage to your blood glucose levels. Make sure that you talk to your doctor regularly. This is particularly important for children and adolescents or if you are an older person.
- Your doctor will also check at least once a year how well your kidneys work. You may need more frequent checks if you are an older person or if your kidneys are not working normally.

### **How to take Glucophage**

Take Glucophage with or after a meal. This will avoid you having side effects affecting your digestion.

Do not crush or chew the tablets. Swallow each tablet with a glass of water.

- If you take one dose a day, take it in the morning (breakfast)
- If you take two divided doses a day, take them in the morning (breakfast) and evening (dinner)
- If you take three divided doses a day, take them in the morning (breakfast), at noon (lunch) and in the evening (dinner)

If, after some time, you think that the effect of Glucophage is too strong or too weak, talk to your doctor or pharmacist.

### **If you take more Glucophage than you should**

If you have taken more Glucophage than you should have, you may experience lactic acidosis.

Symptoms of lactic acidosis are vomiting, bellyache (abdominal pain) with muscle cramps, a general feeling of not being well with severe tiredness, and difficulty in breathing. If this happens to you, you may need immediate hospital treatment, as lactic acidosis may lead to coma. Contact a doctor or the nearest hospital straight away.

### **If you forget to take Glucophage**

Do not take a double dose to make up for a forgotten dose. Take the next dose at the usual time.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

## **4. POSSIBLE SIDE EFFECTS**

Like all medicines, Glucophage can cause side effects, although not everybody gets them. The following side effects may occur:

### **Very common side effects (in more than 1 in 10 people)**

- digestive problems, such as feeling sick (nausea), being sick (vomiting), diarrhoea, bellyache (abdominal pain) and loss of appetite. These side effects most often happen at the beginning of the treatment with Glucophage. It helps if you spread the doses over the day and if you take Glucophage with or straight after a meal. **If symptoms continue, stop taking Glucophage and talk to your doctor.**

### **Common side effects (in less than 1 in 10 people)**

- changes in taste.

### **Very rare side effects (in less than 1 in 10,000 people)**

- lactic acidosis. This is a very rare but serious complication particularly if your kidneys are not working properly. Symptoms of lactic acidosis are vomiting, bellyache (abdominal pain) with muscle cramps, a general feeling of not being well with severe tiredness, and difficulty in breathing. **If this happens to you, you may need immediate hospital treatment, as lactic acidosis may lead to coma. Stop taking Glucophage immediately and contact a doctor or the nearest hospital straight away.**
- abnormalities in liver function tests or hepatitis (inflammation of the liver; this may cause tiredness, loss of appetite, weight loss, with or without yellowing of the skin or whites of the eyes). If this happens to you, **stop taking Glucophage and talk to your doctor.**
- skin reactions such as redness of the skin (erythema), itching or an itchy rash (hives).
- low vitamin B12 levels in the blood.

### **Children and adolescents**

Limited data in children and adolescents showed that adverse events were similar in nature and severity to those reported in adults.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE GLUCOPHAGE**

Keep out of the reach and sight of children. If a child is treated with Glucophage, parents and caregivers are advised to oversee how this medicine is used.

This medicinal product does not require any special storage conditions.

Do not use Glucophage after the expiry date which is stated on the carton or the bottle or the blister after 'EXP'. The expiry date refers to the last day of that month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## **6. FURTHER INFORMATION**

### **What Glucophage contains**

- The active substance is metformin hydrochloride. One film-coated tablet of Glucophage 1000 mg contains 1000 mg metformin hydrochloride corresponding to 780 mg metformin base.
- The other ingredients are povidone K 30, magnesium stearate, hypromellose, macrogol 400 and macrogol 8000.

### **What Glucophage looks like and contents of the pack**

Glucophage 1000 mg film-coated tablets are white, oval and biconvex with a score line on both sides and with '1000' engraved on one side. The tablet can be divided into equal halves.

The tablets are supplied in blister packs of 1 (x30), 20, 30, 50, 60, 90, 100, 120, 180 or 600 tablets or in plastic bottles with child-resistant caps of 20, 30, 50, 60, 90, 100, 120, 180 or 600 tablets.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

#### **Marketing Authorisation Holder**

[to be completed nationally]

#### **Manufacturer**

Merck Santé s.a.s.  
2 rue du Pressoir Vert  
45400 Semoy  
France

or

Merck KGaA  
Frankfurter Str. 250  
64293 Darmstadt  
Germany

or

Merck KGaA & Co. Werk Spittal  
Hösslgasse 20  
9800 Spittal/Drau  
Austria

or

Merck S.L.  
Poligono Merck  
Mollet Del Vallès 08100 Barcelona  
Spain

or

N. Petsiavas S.A.  
21, Agion Anargyron St., Kalyftaki, Kato Kifissia  
Athens 145 64  
Greece

or

Famar Lyon  
29 avenue Charles de Gaulle  
69230 Saint-Genis Laval  
France

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Glucophage: Austria, Belgium, Bulgaria, Cyprus, Czech Republic, Finland, France, Germany, Greece, Iceland, Ireland, Italy, Latvia, Luxembourg, Malta, Norway, Poland, Romania, Slovak Republic, Slovenia, Sweden

Merckformin: Hungary

Risidon: Portugal

**This leaflet was last approved in {MM/YYYY}.**

[To be completed nationally]